




Accurate Solution of the Hartree-Fock Equation by the Tensor-structured Methods

Venera Khoromskaia

Scientific Computing Group, MIS MPI, Leipzig



Two-level Tucker-canonical tensor format in \mathbb{R}^d

We approximate the d -th order tensors (function of discrete variables)

$$A = [a_{i_1 \dots i_d}] \in \mathbb{R}^{n^d}, \quad i_\ell = 1, \dots, n, \quad \ell = 1, \dots, d, \quad n \in \mathbb{N}; \quad S(A) = n^d,$$

by the short-term sum of rank-1 tensors.


Canonical model: using a sum of normalized rank-1 tensors

$$A_{(R)} = \sum_{k=1}^R c_k u_k^{(1)} \otimes \dots \otimes u_k^{(d)} \approx A.$$

Tucker model: using orthonormalized set $\{V_{k_\ell}^{(\ell)}\} \in \mathbb{R}^n, \quad k_\ell = 1, \dots, r$,

$$A_{(r)} = \sum_{k_1=1}^r \dots \sum_{k_d=1}^r b_{k_1 \dots k_d} V_{k_1}^{(1)} \otimes \dots \otimes V_{k_d}^{(d)} \approx A.$$

Mixed format:



Tensor Approximation of the Hartree-Fock equation

The Hartree-Fock equation for the N -electron system is a self-consistent eigenvalue problem (EVP)

$$\left(-\frac{\Delta}{2} + V_{nuc} + V_H\right) \varphi_i(\mathbf{x}) - \int_{\mathbb{R}^3} \frac{\tau(\mathbf{x}, \mathbf{y}) \varphi_i(\mathbf{y})}{|\mathbf{x} - \mathbf{y}|} d\mathbf{y} = \lambda_i \varphi_i(\mathbf{x})$$

for $i = 1, \dots, N/2$, where N is the number of electrons in a molecule.

Computation of the Hartree potential.

The Hartree potential

$$V_H = \int_{\mathbb{R}^3} \frac{\rho(\mathbf{y})}{|\mathbf{x} - \mathbf{y}|} d\mathbf{y} \equiv \rho * \frac{1}{|\mathbf{x}|}, \quad \rho(\mathbf{x}) = 2\tau(\mathbf{x}, \mathbf{x}), \quad \tau(\mathbf{x}, \mathbf{y}) = \sum_{i=1}^{N/2} \varphi_i(\mathbf{x}) \varphi_i(\mathbf{y}).$$


V_H is computed by the tensor product convolution.

Electron density ρ is represented by the rank- $R_{\rho r}$ canonical tensor F . Newton potential $\frac{1}{|\mathbf{x}|}$ is approximated by the rank- R_N ($R_N \sim 20 \div 30$) canonical tensor G obtained by the optimised *sinc*-quadratures.

$$V_H \approx F * G = \sum_{k=1}^{R_N} \sum_{m=1}^{R_{\rho r}} c_k b_m (u_1^k * v_1^m) \otimes (u_2^k * v_2^m) \otimes (u_3^k * v_3^m)$$

which leads to the cost $O(RR_1 n \log n) \ll O(n^3 \log n)$, where the latter corresponds to 3D FFT.

The initial large rank $R_\rho \sim R_0^2 \sim 10^4$ is reduced to $R_{\rho r} \sim 10^2$ by the **Multigrid Accelerated canonical-to-Tucker algorithm (C2T)**.





Electron density of the water molecule (left) and the times for computation of the C2T and 3D convolution (right), up to the grid size 16384^3 (in Matlab).

Computation of the Coloumb matrix.



$$J_{km} := \int_{\mathbb{R}^3} g_k(\mathbf{x}) g_m(\mathbf{x}) V_H(\mathbf{x}) d\mathbf{x}, \quad k, m = 1, \dots, R_0,$$

where $\{g_k\}$ represents the GTO basis set.



MG versus unigrid C2T times and the abs. error for the Coulomb matrix of H_2O . **Computation of the Exchange matrix.**

$$\{K_{ex}\}_{k,m} := -\frac{1}{2} \int_{\mathbb{R}^3} g_k(\mathbf{x}) \frac{\tau(\mathbf{x}, \mathbf{y})}{|\mathbf{x} - \mathbf{y}|} g_m(\mathbf{y}) d\mathbf{x} d\mathbf{y}, \quad k, m = 1, \dots, R_0$$





Abs. error for K_{ex} of CH_4 ($\sim 10^{-4}$) and the pseudopotential of CH_3OH ($\sim 10^{-5}$).

Iterative solution of the EVP in tensor format

Discrete Hartree Fock equation in GTO basis with the Fock matrix $F = H_0 + J(\rho(x)) - K_{ex}(\tau(x, y))$.

Our discrete version of the SCF method is based on:

- Initial guess $J = 0, K = 0$.
- Computation on a sequence of refined grids $n = n_0, 2n_0, \dots, 2^p n_0$.
- Solving EVP $[H_0 + J(\rho_m(x_n)) - K_{ex}(\tau_m(x_n, y_n))] \phi_{i,m} = \lambda_{i,m} \phi_{i,m}$
- Fast update of $J(\rho(x_n))$ and $K_{ex}(\tau(x_n, y_n))$ by computations in



Convergence of the eigenvalues in the EVP (left) w.r.t. iterations and of the E_{HF} energy w.r.t the univariate grid size of the EVP (right).

$$E_{HF} = 2 \sum_{i=1}^{N/2} \lambda_i - \sum_{i=1}^{N/2} (\tilde{J}_i - \tilde{K}_i), \quad \tilde{J}_i = (\psi_i, V_H \psi_i)_{L^2}, \quad \tilde{K}_i = (\psi_i, V_{ex} \psi_i)_{L^2}.$$

References

- H.-J. Flad, V. Khoromskaia and B. N. Khoromskij. Solution of the Hartree-Fock Equation in the Tensor-product Format. in progress, 2009.
- V. Khoromskaia. Computation of the Exchange matrix by Tensor product Convolution on 3D cartesian Grids. in progress, 2009.
- V. Khoromskaia and B. N. Khoromskij. Multigrid Accelerated Tensor Approximation of Function Related Multi-dimensional Arrays. Preprint 40/2008, MPI MIS Leipzig, 2008. SIAM J. Scientific Computing, 2009, (accepted).
- B. N. Khoromskij and V. Khoromskaia, Low Rank Tucker Tensor Approximation to the Classical Potentials. Centr. European J. of Math., 5(3) (2007), 1-28.
- B. N. Khoromskij, S.R. Chinnamsetty, H.-J. Flad and V. Khoromskaia. Tensor Decomposition in Electronic Structure Calculations on 3D Cartesian Grids. Preprint 65, MPI MIS Leipzig, 2007; JCP, (submitted).